




Nutrition Situation Update Report 2018

SOURCE: OPT PLUS REPORT 2017
National Nutrition Council (NNC), Region VI




TREND ON THE PREVALENCE OF UNDERNUTRITION AMONG PRE-SCHOOL CHILDREN (PSC) REGION VI, 2013-2017







PERCENT DISTIBUTION OF UNDERNUTRITION (Weight for Age) AMONG PSC ACCORDING TO NUTRITIONAL STATUS REGION VI, 2017



SUW: 8,811

■ UW: 36,605

NORMAL: 916,908

■ AN: 10,003


ACTUAL PSW: 972,327

ESTIMATED PS: 1,272,234


%COVERAGE: 76.43%



RANKING OF PROVINCES ACCORDING TO PREVALENCE OF UNDERNUTRITION (Weight for Age) AMONG PSC REGION VI, 2017




RANKING OF PROVINCES ACCORDING TO PERCENT COVERAGE OF UNDERNUTRITION (Weight for Age) AMONG PSC WEIGHED REGION VI, 2017




*All Provinces have not reached 80% coverage



RANKING OF CITIES ACCORDING TO PREVALENCE OF UNDERNUTRITION (Weight for Age) AMONG PSC REGION VI, 2017




RANKING OF CITIES ACCORDING TO PERCENT COVERAGE OF UNDERNUTRITION (Weight for Age) AMONG PSC WEIGHED REGION VI, 2017






TOP 10 NUTRITIONALLY DEPRESSED MUNICIPALITIES ACCORDING TO PREVALENCE OF UNDERNUTRITION (Weight for Age) with 80 % COVERAGE AMONG PSC REGION VI, 2017





PERCENT DISTRIBUTION OF STUNTING (Length/Height for age) AMONG PSC ACCORDING TO NUTRITIONAL STATUS REGION VI, 2017



Normal: 853,040

Stunted: 77,481

Severely Stunted: 27,545

■ Tall: 14,261


ACTUAL PSW: 972,327

ESTIMATED PS: 1,272,235

%COVERAGE: 76.43%




RANKING OF PROVINCES ACCORDING TO **PREVALENCE OF STUNTING (Length/Height for age) AMONG PSC REGION VI, 2017**






RANKING OF CITIES ACCORDING TO PREVALENCE OF STUNTING (Length/Height for age) AMONG PSC REGION VI, 2017






TOP 10 MUNICIPALITIES ACCORDING TO PREVALENCE OF STUNTING (Length/Height for age) WITH 80% COVERAGE AMONG PSC **REGION VI, 2017**





PERCENT DISTRIBUTION OF WASTING (Weight for Length) AMONG PSC ACCORDING TO NUTRITIONAL STATUS REGION VI, 2017



Normal: 820,804

■ Wasted: 21,712

Severely Wasted: 8,440

Overweight: 14,704

Obese: 9,902


Actual PSM: 875,562

Estimated PS: 1, 235, 466

☐ % coverage: 70.87%




RANKING OF PROVINCES ACCORDING TO **PREVALENCE OF WASTING (Weight for Length) AMONG PSC REGION VI, 2017**






RANKING OF CITIES ACCORDING TO PREVALENCE OF WASTING (Weight for Length) AMONG PSC REGION VI, 2017





TOP 10 MUNICIPALITIES ACCORDING TO PREVALENCE OF WASTING (Weight for Length) WITH 80 % COVERAGE AMONG PSC REGION VI, 2017






PREVALENCE OF OVERWEIGHT (WEIGHT FOR HEIGHT /LENGTH) AMONG PSC REGION VI, 2017



Note: Calinog, Iloilo NO data




RANKING OF PROVINCES ACCORDING TO PREVALENCE OF OVERWEIGHT (Weight for Length) AMONG PSC REGION VI, 2017



Calinog, Iloilo NO data



RANKING OF CITIES ACCORDING TO PREVALENCE OF OVERWEIGHT (Weight for Length) AMONG PSC REGION VI, 2017






TOP 10 MUNICIPALITIES ACCORDING TO PREVALENCE OF OVERWEIGHT (Weight for Length) **AMONG PSC IN REGION VI, 2017**






PREVALENCE OF OBESE ACCORDING TO WEIGHT FOR HEIGHT /LENGHT AMONG PSC REGION VI, 2017



Note: Calinog, Iloilo NO data



PREVALENCE OF OBESE (Weight for Length) AMONG PSC BY PROVINCE IN REGION VI, 2017






PREVALENCE OF OBESE (Weight for Length) AMONG PSC BY CITIES IN REGION VI, 2017





TOP 10 MUNICIPALITIES ACCORDING TO PREVALENCE OF OBESE (Weight for Length) AMONG PSC IN REGION VI, 2017







5 out of 100


4.67% are undernourished



3 out of 100

3.44% are wasted





1 out of 10

10.80% are stunted



SUMMARY



2 out of 100

1.68% are overweight



1 out of 100


1.13% are obese



Children, Less than 5 years old

 Prevalence rate of undernutrition, stunting, and wasting generally decreased except overweightfor-height and obesity.

Summary



PLEASE



US ON...



www.facebook.com/nncr6

www.nnc.gov.ph/regional-offices/region-vi



